

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/37, 15/62, C07K 14/025, A61K 39/12		A1	(11) International Publication Number: WO 96/11274
			(43) International Publication Date: 18 April 1996 (18.04.96)
(21) International Application Number: PCT/US95/12914		(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 6 October 1995 (06.10.95)		2201601 Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 08/319,467 6 October 1994 (06.10.94) US			
(71) Applicant: THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US).			
(72) Inventors: LOWY, Douglas, R.; 4709 South Chelsea Lane, Bethesda, MD 20814 (US). SCHILLER, John, T.; 11306 Maple View Drive, Silver Spring, MD 20902 (US). GREENSTONE, Heather; 1901 Stratton Road, Silver Spring, MD 20910 (US).			
(74) Agent: ALTMAN, Daniel, E.; Knobbe, Martens, Olson & Bear, Suite 1600, 620 Newport Center Drive, Newport Beach, CA 92660 (US).			
(54) Title: CHIMERIC PAPILLOMAVIRUS-LIKE PARTICLES			
(57) Abstract The present invention provides a papillomavirus-like particle, characterized as having conformational epitopes, comprising a papillomavirus L1 product and a papillomavirus L2 fusion product; and related synthetic DNA molecules, host cells, methods and vaccines.			

CHIMERIC PAPILLOMAVIRUS-LIKE PARTICLES**FIELD OF THE INVENTION**

The present invention relates to chimeric papillomavirus-like particles and related synthetic DNA molecules, host cells, methods and vaccines.

BACKGROUND OF THE INVENTION

Papillomaviruses infect the epithelia of humans and a wide variety of animals, where they generally induce benign proliferation at the site of infection. However, in some cases the lesions induced by certain papillomaviruses undergo malignant progression. There is a strong association between malignant progression of human genital lesions and certain human papillomavirus (HPV) types, such as HPV16. Infection by one of these types is considered the most significant risk factor in the development of cervical cancer, one of the most common cancers of women worldwide (zur Hausen, H., Science 254:1167 (1991); Schiffman, M.H., J. Natl. Cancer Inst. 84:394 (1992)). The majority of cervical carcinomas contain and express HPV early genes, such as E6 and E7, and these genes have been shown to have potent transforming and immortalizing activity in cultured cells (Werness, B.A., Munger, K. & Howley, P.M. (1991) Advances in Oncology, eds. DeVita V.T., Hellman, S. & Rosenberg, S.A. (Lippincott, Philadelphia) pp.3-18).

Papillomaviruses are non-enveloped double-stranded DNA viruses about 55 nm in diameter with an approximately 8 kb genome in the nucleohistone core (Baker, et al, Biophys J 60:1445 (1991)). The capsids are composed of two virally-encoded proteins, L1 and L2, that migrate on SDS-PAGE gels at approximately 55 kDa and 75 kDa, respectively (Mose Larson et al., J. Virol. 61:3596 (1987)). L1, which is the major capsid protein, is arranged in 72 pentameters which associate with T-7 icosahedral symmetry. The function and position within the virion of L2 are unclear (Baker, et al, Biophys J 60:1445 (1991)).

The L1 protein has the capacity to self-assemble so that large amounts of virus-like particles (VLPs) may be generated by expression of the L1 protein from a number of species of papillomavirus in a variety of recombinant expression systems (Hagensee et al., J Virol 67:315 (1993); Kirnbauer et al., Proc Natl Acad Sci USA 89:12180 (1992); Kirnbauer et al., J Virol 67:6928 (1993); Rose et al., J Virol 67:1936 (1993)). Although not required for assembly, L2 is incorporated into VLPs when co-expressed with L1 (L1/L2 VLPs) in cells.

Immunization of rabbits with native virions or L1 VLPs, but not with non-assembled L1 expressed in *E. coli*, induces high titers of neutralizing serum antibodies (Christensen, N.D. and Kreider, J.W., J Virol 64:3151 (1990); Kirnbauer et al., Proc Natl Acad Sci USA 89:12180 (1992); Pilacinski et al., Bio/Technology 2:356 (1984); Segre et al., Am. J. Vet. Res. 16:517 (1955)). The polyclonal and monoclonal antibodies generated against native particles recognize conformationally dependent epitopes (Christensen, N.D. and Kreider, J.W., Virus Res 28:195 (1993); Christensen et al., J Virol 64:5678 (1990); Christensen et al., Virology 181:572 (1991)).

Neutralizing antibodies generated against VLPs also recognize conformationally dependent epitopes. Using infectious BPV1, which can be readily obtained from bovine lesions, and a quantitative *in vitro* BPV1 infectivity assay (Dvoretzky et al., Virology 103:369 (1980)), workers showed VLPs from bovine papillomavirus induced high levels of neutralizing antibodies (Kirnbauer et al., Proc Natl Acad Sci USA 89:12180 (1992)). The neutralizing antibodies

- 2 -

were directed against conformationally dependent epitopes, in that denaturation of the particles prior to immunization abolished the ability of the preparation to induce
5 neutralizing activity (*id.*).

When the L1 gene of a HPV16 isolate derived from a nonprogressed lesion was used to express the L1 major capsid protein in insect cells via recombinant baculoviruses, L1 self-assembled into VLPs at a yield 3 orders of magnitude higher than what had been obtained using L1 derived from the prototype HPV16 (originally isolated from a cancerous
10 lesion), and formed VLPs that were morphologically similar to native virions (Kirnbauer et al., J Virol 67:6929 (1993)). DNA sequence comparison identified a single nonconserved amino acid change to be responsible for the inefficient self-assembly of the prototype L1 (*id.*). The L1 of the assembly-competent clone is thus considered to be the wild-type gene, and the prototype L1 of the assembly-defective clone a mutant.

15 Using HPV16 VLPs of the wild-type L1 protein as antigens, an ELISA was developed that detected serum antibodies in patients infected with HPV16 (Kirnbauer et al., J. Natl. Cancer Inst. 86:494 (1994)). In contrast, neither denatured HPV16 particles nor preparations of the prototype L1 protein could detect these antibodies (*id.*). These results demonstrate that the prototype L1 protein does not present conformational epitopes.

20 Rabbit serum raised against self-assembled wild-type HPV16 L1/L2 virus-like particles was discovered to prevent HPV16 VLP binding to cell surface molecules (Roden et al., J. Virol., 68:7260-7266 (1994)). In contrast, serum raised against the prototype strain of HPV16 L1/L2 did not prevent such binding (*id.*). The data show that the prototype HPV16 strain lacks conformational epitopes.

25 Rabbits immunized with intact cottontail rabbit papillomavirus (CRPV) virus-like particles composed of L1 or L1/L2 were protected from subsequent experimental challenge by infectious CRPV (Breitburd et al., J. Virol. 69:3959-3963 (June, 1995)). In contrast, those immunized with denatured particles were not protected (*id.*). These findings are consistent with the conclusion that VLPs presenting conformational epitopes are able to
30 induce protective immunity.

- 2a -

VLPs composed of capsid proteins are attractive candidates for prophylactic vaccines to prevent papillomavirus infection. However, it is unlikely that these VLP
5 vaccines would have therapeutic effects against established papillomavirus infections. The capsid proteins, unlike E6 and E7, are not detectably expressed in progressed lesions or in infected basal epithelial cells, which are the presumed targets in immune regression of papillomas.

There is evidence from experimental models that immunity against papillomavirus
10 proteins other than L1 and L2 might help control papillomavirus infection. Since E6 and E7 are selectively maintained during oncogenic progression, there is the possibility that peptides derived from these oncoproteins could serve as targets for cell-mediated immune responses to HPV-containing tumor cells. Studies in animal models suggest the E7 protein of HPV16 acts as a tumor rejection antigen (Chen et al., Proc. Natl. Acad. Sci. USA
15 88:110 (1991); Feltkamp et al., Eur. J. Immunol. 23:2242 (1993)). Moreover, the frequency of HPV infection, persistence of HPV infection, and risk of developing cervical cancer and other HPV-related cancers is increased in patients with depressed cellular immunity (Allout et al., Br. Med. J. 298:153 (1989); Laga et al., Int. J. Cancer 50:45 (1992)). These observations suggest

cell-mediated immunity is important in the defense against HPV infection and its associated tumor development. The induction of such immunity might be therapeutic, as well as prophylactic.

It has been demonstrated that foreign peptides can be incorporated into viral capsid-like structures and these chimeric particles can be used to present foreign antigens to the immune system. Published examples include hepatitis B core antigen particle presentation of human rhinovirus type 2 epitopes (Francis et al., Proc. Natl. Acad. Sci. USA 87:2545 (1990)), gp41 of HIV (Borisova et al., FEBS Lett. 259:121 (1989)), and B19 parvovirus particle presentation of peptides from herpes simplex virus 1 and murine hepatitis virus (Brown et al., Virology 198:477 (1994)). The parvovirus chimeras protected mice from experimental challenge with the corresponding virus. In all of the above systems, foreign sequences have been inserted in proteins integral to the capsid structure and have been limited to less than 20 amino acids. However a recent study (Miyamura et al., Proc. Natl. Acad. Sci. USA 91:8507 (1994)) has demonstrated that the entire 147 aa hen egg white lysozyme protein can be incorporated into B19 parvovirus particles when fused to the parvovirus L1 minor capsid protein. The lysozyme remained biologically active and elicited an immune response when injected into rabbits. In perhaps less relevant studies, hepatitis B virus surface antigen particles (which are lipid membrane structures) containing 84 aa of HIV-1 envelope glycoprotein (Michel et al., J. Virol. 64:2452 (1990)) and yeast Ty virus-like particles containing a portion of HIV-1 V3 loop (Griffiths et al., J. Virol. 65:450 (1991)) have also been shown to produce an immune response to the inserted peptides when inoculated into animals. With respect to papillomaviruses, it was recently reported that hepatitis B core antigen particles containing HPV16 E7 peptides (all less than 20 aa) induced peptide specific antibodies and T-helper responses in mice (Tindle et al., Virology 200:547 (1994)).

Chimeric particles based on self-assembled papillomavirus L1 have not been reported, nor has the use of L2 as a viral fusion partner for purposes of generating chimeric VLPs been described. The chimeric particle studies cited above involve viruses that are unrelated to papillomaviruses and thus cannot predict the results of chimeric particle studies involving papillomaviruses. Indeed, in the papillomavirus study by Kimbauer et al., J Virol 67:6929 (1993), *supra*, it was demonstrated that a single nonconserved amino acid change in L1 is responsible for efficient self-assembly of L1 into VLPs and the presentation of conformational epitopes, which seem to be required for induction and detection of clinically relevant immune reactivity. Thus, the studies using viruses unrelated to papillomavirus cannot predict whether a papillomavirus L2 containing a foreign peptide or protein can co-assemble with papillomavirus L1 into particles, given that a single amino acid substitution in L1 can abolish efficient self-assembly. Neither can these studies predict whether any resulting chimeric particles will retain the ability to induce or detect neutralizing antibodies or other immune related responses, given that a single amino acid substitution in L1 can bar the presentation of conformational epitopes.

It is an object of the present invention to provide chimeric papillomavirus-like particles. These chimeric particles may function as platforms for multivalent antigen presentation. Or they may serve for delivery into cells of proteins for processing into peptides and subsequent presentation of these peptides within the context of MHC molecules to elicit a cell-mediated immune response. The chimeric particles represent a cost effective way to generate an effective papillomavirus vaccine with a broad spectrum of utility. Alternatively, the chimeric

papillomavirus-like particles may be applied to VLP and/or fusion partner purification. Or, the particles may operate to deliver into cells intact and active proteins, for example, enzymes, or toxins or drugs.

SUMMARY OF INVENTION

10 This invention provides a papillomavirus-like particle having conformational epitopes comprising a papillomavirus L1 fusion product. This invention also provides a composition comprising the aforementioned papillomavirus-like particle.

15 This invention also provides a composition comprising a first synthetic DNA molecule characterized as encoding a papillomavirus L1 product and a second synthetic DNA molecule characterized as encoding a papillomavirus L2 fusion product, wherein said molecules are capable of directing expression in a transformed cell of papillomavirus-like particles characterized as having conformational epitopes, wherein said particles comprise said papillomavirus L1 product and papillomavirus L2 fusion product.

20

25

- 4a -

According to one aspect of the invention, there is provided a papillomavirus-like particle, characterized as having conformational epitopes, comprising a papillomavirus L1
5 product and a papillomavirus L2 fusion product.

The papillomavirus L2 fusion product may be characterized as being a human papillomavirus L2 fusion product or a bovine papillomavirus L2 fusion product.

The human papillomavirus L2 fusion product may be characterized as being a HPV16 L2 fusion product and the bovine papillomavirus L2 fusion product may be
10 characterized as being a BPV1 L2 fusion product.

The papillomavirus L2 fusion product may comprise a fusion partner characterized as being a peptide or a full-length protein, or may comprise fusion partners that include peptides or full-length proteins or combinations thereof linked in tandem.

The fusion partner may be a papillomavirus E6 or E7 product.

15 The papillomavirus L2 fusion product may be characterized as being fused at its 5' or 3' end to a fusion partner.

The papillomavirus L2 fusion product may be characterized as having a fusion partner inserted between L2 amino acids.

The papillomavirus L1 product may be characterized as being a human
20 papillomavirus L1 product or a bovine papillomavirus L1 product.

The human papillomavirus L1 product may be characterized as being a HPV16 L1 product and the bovine papillomavirus L1 product may be characterized as being a BPV1 L1 product.

The papillomavirus-like particle may consist essentially of HPV16L1 and
25 HPV16L2-HPV16E7 (full-length), where HPV16E7 is fused to the 3' end of HPV16L2, BPVL1 and BPVL2-HPV16E7 (full-length), where HPV16E7 is fused to the 3' end of BPVL2, or BPVL1 and BPVL2-HPV16E7 (amino acids 1-30), where HPV16E7 is fused to BPVL2 between L2 amino acids 274 and 275.

- 4b -

According to another aspect of the invention, there is provided one or more synthetic DNA molecule or molecules characterized as singly or doubly encoding a papillomavirus L1 product and a papillomavirus L2 fusion product where the molecule or
5 molecules direct expression in a transformed host cell of a papillomavirus-like particle, characterized as having conformational epitopes, comprising the papillomavirus L1 product and the papillomavirus L2 fusion product.

The transformed host cell may be an insect host cell (such as Sf9 insect host cell)
10 and the DNA molecule or molecules may further comprise an insect cell vector (such as a baculovirus vector), or the transformed host cell may be a mammalian host cell and the DNA molecule or molecules may further comprise a mammalian cell vector (such as a vaccinia virus vector), or the transformed host cell may be a yeast host cell and the DNA molecule or molecules may further comprise a yeast cell vector.

According to another aspect of the invention, there is provided a host cell transformed with the DNA molecule or molecules of above.

According to yet another aspect of the invention, there is provided a method for using the DNA molecule or molecules of above comprising the steps of: providing conditions for the molecule or molecules of above to direct the above expression; and recovering the papillomavirus-like particle from the above transformed host cell.

According to still another aspect of the invention, there is provided a method for producing a papillomavirus-like particle, characterized as having conformational epitopes, comprising a papillomavirus L1 product and a papillomavirus L2 fusion product, which method comprises the step of providing conditions for the DNA molecule or molecules of above to direct the above expression in the above transformed host cell of the papillomavirus-like particle.

The invention also provides a method of purification of the papillomavirus-like particle of above comprising the step of exposing the papillomavirus-like particle to an affinity chromatography column, comprising antibodies that bind to a fusion partner of the papillomavirus L2 fusion product of the papillomavirus-like particle, resulting in the purification of the particle.

The invention further provides a method of purification of a fusion partner of the papillomavirus L2 fusion product of the papillomavirus-like particle of above comprising the step of isolating the papillomavirus-like particle of above resulting in the purification of the fusion partner.

The invention additionally provides a method of delivery into a cell of a fusion partner of the papillomavirus L2 fusion product of the papillomavirus-like particle of above comprising the step of administering the papillomavirus-like particle of above to the cell resulting in the delivery into the cell of the fusion partner.

The invention moreover provides a vaccine comprising the papillomavirus-like particle of above.

DETAILED DESCRIPTION OF THE INVENTION

This invention arises from the result that a papillomavirus L2 fusion product can become incorporated into a papillomavirus L1 product-based papillomavirus-like particle that presents conformational epitopes.

This result was unexpected.

In the papillomavirus study by Kimbauer et al., J Virol 67:6929 (1993), *supra*, it was demonstrated that a single nonconserved amino acid change in L1 is responsible for efficient self-assembly of L1 into VLPs and the presentation of conformational epitopes, which seem to be required for induction and detection of clinically relevant immune reactivity. Thus, it could not be predicted whether a papillomavirus L2 containing a foreign peptide or protein could co-assemble with papillomavirus L1 into particles, given that a single amino acid substitution in L1 can abolish efficient self-assembly. Neither could it be predicted whether any resulting chimeric particles would retain the ability to induce or detect neutralizing antibodies or other immune related responses, given that a single amino acid substitution in L1 can bar the presentation of conformational epitopes.

A papillomavirus L2 fusion product is meant to include a chain of amino acids in which part of the chain comes from a L2 protein sequence and part of the chain comes from another protein sequence (or other protein sequences).

L2 fusion products are produced by splicing (in frame) an open reading frame for one protein (or a number of proteins) next to or into an open reading frame for L2.

Protein engineering is used to determine the structure of the L2 fusion product. In a routine exercise for protein engineers, they generate variants of the natural protein L2. The changes they make can be educated guesses based on detailed knowledge of the structure of L2; alternatively, changes can be made on a purely random basis. Or a combination of structural information with random mutagenesis and selection can have dramatic results.

For example, the selection of L2 amino acids between which to insert a fusion peptide or protein can be made on this basis. The existence of a region where amino acid sequence and length vary between papillomaviruses suggests that this region represents a structure that is nonessential for the integrity of L2 and/or its incorporation into particles. The observed ability to insert a fusion peptide or protein within L2 implies that such a region exists.

Determination of a structure for the L2 fusion product is performed on the basis of the ability of the L2 fusion product to become incorporated into a papillomavirus L1 product-based papillomavirus-like particle that presents conformational epitopes.

Preferred means of determining a structure include assays that measure the efficiency and/or authenticity of incorporation of an L2 fusion product into a papillomavirus L1 product-based papillomavirus-like particle.

The efficiency and/or authenticity of formation of L1/L2 VLPs is related to the presentation of conformational epitopes (*supra*).

The chimeric L2 will thus become incorporated into L1-based VLPs with an efficiency and/or authenticity similar to that of incorporation of wild-type L2.

Other preferred means of determining a structure for the L2 fusion product include assays that measure the induction and/or detection of neutralizing antibodies.

The induction and/or detection of neutralizing antibodies is related to the presentation of conformational epitopes, because neutralizing antibodies are directed against conformationally dependent epitopes (*supra*).

The incorporation of the chimeric L2 will thus not inhibit the induction and/or detection of neutralizing antibodies by chimeric papillomavirus-like particles as compared to L1 or L1/L2 VLPs.

For purposes of the invention, a chimeric L2 fusion product is capable of becoming incorporated into a papillomavirus L1 product-based papillomavirus-like particle that presents conformational epitopes, whether identified by the above means or any other means known in the art.

The structure of L2 and its fusion partner (and L1) is meant to include a full-length L2 protein (and L1 protein) and a full-length fusion partner protein, and their peptide fragments, whether 5' fragments, 3' fragments, or internal fragments, having at least about 20 amino acid residues, advantageously at least about 10 amino acid residues, and preferably at least about 5 amino acid residues. Type, subgroup and strain variations of L2 (and L1), and human allelic and species variations of the fusion partner protein, are expressly contemplated as falling within the scope of the invention. The invention also includes conservative variants of the full-length L2 protein (and L1 protein) and the full-length fusion partner protein, and their peptide fragments, where conservative amino acids are substituted for amino acid residues of wild-type L2 (and L1) and the fusion partner protein. To account for

degeneracy of the genetic code, the invention also includes DNA coding for the same amino acid residues as does the DNA of the L2 (and L1) and fusion partner gene.

The chimeric papillomavirus-like particle itself is envisioned as incorporating any L2 fusion product with any L1 product from any papillomavirus, whether the genomes are closely related, or are distantly related, so long as incorporation into particles occurs. Thus, a L2 fusion product, for example, related to any of BPV-1, BPV-2, BPV-4, CRPV, DPV, EEPV, HPV-1, HPV-5, HPV-6, HPV-8, HPV-11, HPV-16, HPV-18, HPV-31 or HPV-33, can be incorporated into particles with any L1 product, for example, from any of the above virus, or any type, subgroup or strain variation of papillomavirus.

Because VLPs present conformationally dependent epitopes required for the induction of high titer neutralizing serum antibodies, VLP chimeras containing L2 fusion products can operate prophylactically as optimized subunit vaccines for the stimulation of humoral immunity to prevent papillomavirus infection and thereby preclude the development of papillomavirus associated cancers and other papillomavirus associated pathologies.

Because it is unlikely VLPs will prove effective as therapeutic vaccines to induce regression of existing papillomavirus proliferative lesions, as discussed above, chimeric VLPs can function to address this long-felt and heretofore unsatisfied need to develop a therapeutic vaccine.

Chimeric VLPs are expected to bind specific cell surface receptors, get internalized and be released into the cytoplasm, and thus be more likely to promote the presentation of peptides in conjunction with class I MHC molecules for display to cytotoxic T cells for the generation of cell-mediated immunity. This is in contrast to uncomplexed proteins that would not be expected to specifically enter cells, or to promote the presentation of peptides in the context of Class I MHC molecules to elicit a cytotoxic T cell response (being more likely, if at all, to promote the presentation of peptides to be linked to Class II MHC molecules and displayed to helper T cells).

Inclusion of one or more fusion partners as L2 fusion products would clearly be a cost effective way to generate an effective papillomavirus vaccine with a broad spectrum of utility, including therapy and improved prevention of clinical lesions.

The fusion partner may be selected from the list consisting of those fusion partners that would provide a method for expanding the potential targets of a VLP-based vaccine, for example, E6 or E7 peptide or full-length E6 or E7, other papillomavirus peptides or proteins, or peptides or proteins of other STD or infectious agents, *e.g.*, Herpes simplex, HIV, Chlamydia trachomatis, Neisseria gonorrhoeae, and Treponema pallidum.

The L2 fusion product is not limited to a single fusion partner per L2 molecule, and may include additional fusion partners, for example, additional peptides or full-length proteins or combinations thereof derived from the same or different proteins linked in tandem. Also, more than one L2 fusion product may be co-assembled into a single VLP. For example, a L2 fusion product or chimeric VLP containing a viral target epitope may also be engineered to contain a binding domain of a co-stimulatory protein or an accessory receptor or ligand involved in immune reactivity, *e.g.*, B7 (which interacts with CD28 on T cells), an intercellular adhesion molecule (ICAM), a lymphocyte functional antigen (LFA), a vascular cell adhesion molecule (VCAM-1), and a heat stable antigen (HSA). (To target the co-stimulators or ligands to the surface of VLPs, see Example 14.)

It will be appreciated that the actual preferred amounts of chimeric papillomavirus-like particles as vaccines in the prevention and/or treatment of disease will vary according to the specific compositions formulated, the mode of application, the particular situs and the organism being treated. Dosages for a given host can be determined using conventional considerations, *e.g.*, by means of an appropriate, conventional vaccination protocol.

5 Alternatively, chimeric VLPs can be used in a method of purification of VLPs. VLPs are useful *per se* as prophylactic vaccines and in immunodiagnostics (*supra*). Briefly, antibodies to the fusion partner are generated using standard immunological techniques, an affinity chromatography column is constructed using the antibodies, and the VLPs are subsequently purified in an affinity chromatography step.

10 Or, chimeric VLPs can be used in a method of purification of a fusion partner. For example, VLPs containing a L2-E7 fusion product may be useful as a means of obtaining correctly folded E7. It has been reported that a greater percentage of cervical cancer patients are seropositive for conformationally correct E7 than for denatured E7, as isolated from bacteria (Viscidi et al., *Int. J. Cancer* 55:780 (1993)). The *in vitro* transcription-translation system used to generate E7 in this report is laborious and expensive, and uses radio-labeled E7. It would be advantageous to purify chimeric E7 VLPs and use the material in an E7 ELISA. To avoid complications of
15 seroreactivity with human L1 or L2, BPV-based particles could be used. Monitoring E7 seroreactivity, which correlates with cancer as opposed to premalignant disease, has been proposed to be useful to follow the course of disease in cervical cancer patients and to screen for reoccurrences (*id.*).

Optionally, chimeric VLPs can be used in a method of delivery of an intact and active protein into cells. This protein may be, for example, an enzyme, or a toxin or a drug. The chimeric VLPs are administered to the cells,
20 either *in vitro*, *in vivo*, *in situ*, or *ex vivo*, and the protein is subsequently delivered into the cell where it functions for its intended purpose, for example, as an enzyme, or a toxin or a drug.

We have succeeded at generating chimeric papillomavirus-like particles incorporating E7 protein or peptide fused to L2.

25 A papillomavirus E7 gene when fused in frame to the 3' end of the L2 gene or within part of the L2 open reading frame expressed an L2-E7 fusion protein that, in the presence of L1 protein, was incorporated into virus-like particles. Efficiency and authenticity of incorporation of L2-E7 into VLPs was similar to that of the wild-type L2 protein. BPV L1/L2-E7 virus-like particles were found to induce neutralizing antibodies as effectively as BPV L1/L2 particles. Chimeric particles were observed to elicit humoral immunity specific for fusion partner epitopes in that rabbits immunized with L2-E7 chimeric particles generated antibodies directed against E7.

30 Having succeeded at generating chimeric papillomavirus-like particles opens the door to fusing virtually any protein or peptide to an L2 product for incorporation into chimeric papillomavirus-like particles.

We accomplished the production of chimeric VLPs by generating L2-E7 fusion proteins and expressing these proteins along with L1 via recombinant baculoviruses in insect cells. We generated three chimeric VLPs: HPV16L1 + HPV16L2-HPV16E7 (full-length), BPV1 + BPV12-HPV16E7 (full-length) and BPV1 + BPV12-HPV16E7 (aa 1-30).

BPV L2 was fused at its 3' end to HPV16 E7. HPV16 L2 was fused at its 3' end to HPV16 E7. Additionally, the first 30 codons of HPV16 E7 were inserted between codons 274 and 275 of BPV L2.

Sucrose gradient co-sedimentation and co-immunoprecipitation of L1 and the L2-E7 chimeras demonstrated the incorporation of the chimeric protein into the VLPs.

5 Transmission electron microscopy revealed that particles containing the L2-E7 fusion protein assembled with the same efficiency as L1 or L1/L2 VLPs, and were morphologically indistinguishable.

In vitro BPV1 neutralization assays demonstrated that the BPV L1-containing chimeric VLPs were capable of inducing neutralizing antisera. Titers were comparable to those obtained using BPV L1/L2 VLPs. Equivalent neutralizing titers of 30,000 were obtained for both BPV L1/L2 VLPs and BPV L1/L2-HPV16E7 (full-length) chimeric VLPs.

10 Rabbits immunized with L2-E7 chimeric particles generated antibodies directed against E7, demonstrating the induction by chimeric VLPs of humoral immunity specific for fusion partner epitopes, and also establishing the location of E7 epitopes as external to VLPs.

As set forth in Example 1, chimeric recombinant baculoviruses can be generated. Genes for the chimeras may be obtained from genomic sources or cDNAs, by direct synthesis, or by any combination thereof. L2 and its fusion partner genes can be amplified by recombinant PCR techniques, for example, with oligos containing restriction enzyme sites and complementarities that facilitate fusion and cloning into expression, transfer and/or cloning vectors, e.g., plasmids.

15 Fused genes can be cloned into a baculovirus expression vector. Another baculovirus expression vector containing L1 can be generated. Or the fused genes can be cloned into a baculovirus double expression vector that already contains L1.

20 Example 2 sets forth a typical procedure for selection of recombinant baculoviruses. CsCl-purified (or equivalent) recombinant plasmid can be cotransfected with baculovirus DNA into Sf9 insect cells by using Lipofectin (or equivalent). The recombinant baculoviruses can then be plaque-purified (for example) using conventional baculovirus vector and insect cell culture procedures.

25 It may be advantageous to use two single expression baculovirus vectors instead of a double expression baculovirus vector. In this case, chimeric VLPs can be produced by infecting Sf9 cells with two recombinant baculoviruses, one encoding a L1 product and the other encoding the L2 fusion product. Locating the genes on different vectors facilitates manipulation of the amount of L1 product and L2 fusion product produced. This approach permits one to change the ratio of L2 fusion product to L1 product in a VLP. Although in native virions, L2 is the minor component as compared to L1, we can achieve greater incorporation of L2 fusion product into VLPs using two single expression vectors.

30 As set forth in Example 3, chimeric particles can be purified by banding in cesium chloride (or equivalent). Sf9 insect cells can be infected, for example, at a multiplicity of infection of 10 with recombinant baculoviruses. After 72 hours (or so), cells can be harvested and sonicated in a phosphate-buffered saline (or equivalent) for 60 sec (or so). After low speed clarification (or equivalent), the lysates can be subjected to centrifugation, for example, at

110,000 x g for 2.5 h through a 40% (wt/vol) sucrose/PBS cushion (SW-28 rotor). The resuspended pellets can be centrifuged to equilibrium, for example, at 141,000 x g for 20 h at room temperature in a 10-40% (wt/wt) CsCl/PBS gradient. The visible band can be harvested, centrifuged to equilibrium again by using the identical conditions, dialyzed extensively against PBS, and stored at 4°C (for example).

5 As set forth in Example 4, co-sedimentation of chimeric complexes can be established, for example, by analytical gradient centrifugation. *E.g.*, a 12 to 45% sucrose step gradient can be allowed to linearize overnight at 4°C, dialyzed samples can be layered on top, and the gradient can be centrifuged at 41,000 rpm (288,000 x g) for 2 h in a SW-41 rotor. Fractions can be harvested and analyzed for co-sedimentation, for example, by Western blotting or co-immunoprecipitation.

10 As set forth in Example 5, co-sedimentation can be established, for example, by co-immunoprecipitation.

 As set forth in Example 6, antisera can be produced. This can be done to conduct a BPV1 neutralization assay (or equivalent) as is described in Dvoretzky et al., Virology 103:369 (1980). *E.g.*, antisera can be produced as follows. Rabbits can be immunized by subcutaneous injection of 330 µl of CsCl gradient-purified particles in PBS at a concentration of 1 mg/ml. Rabbits can then be boosted with the same amount of particles two weeks and four weeks after the primary injection.

15 As set forth in Example 7, a BPV1 neutralization assay (or equivalent) can be performed to test whether BPV chimeric particles present conformational epitopes. *E.g.*, serial dilutions of sera obtained 3 weeks after the second booster injection can be incubated with ≈ 500 focus-forming units of BPV1 virus for 30 min, the virus can be adsorbed to C127 cells for 1 h, and the cells can be cultured for 3 weeks. Foci can then be stained with 0.5% methylene blue/0.25% carbol fuchsin in methanol. Neutralizing titers can be obtained for chimeric VLPs and control BPVL1-L2 VLPs. Equivalent neutralizing titers will indicate proper folding of the chimeric particles effective to present conformational epitopes.

20 As set forth in Example 8, chimeric particles can be assayed, for example, by transmission electron microscopy. *E.g.*, purified particles can be adsorbed to carbon-coated grids, stained with 1% uranyl acetate, and examined with a Philips electron microscope model EM 400T at x36,000 magnification. Efficiency of incorporation and morphology of chimeric particles and L1 or L1/L2 VLPs can be compared and contrasted. Indistinguishable efficiency of incorporation and morphology will indicate proper self-assembly of the chimeric particles.

25 As set forth in Example 9, chimeric particles can be assayed, for example, for induction of humoral immunity specific for fusion partner epitopes. *E.g.*, rabbits can be inoculated with chimeric VLPs. The sera can be analyzed for antibodies, for example, by subjecting a sample of the fusion partner antigen to SDS-PAGE and then Western blotting. Immune and preimmune (control) sera can be applied at an appropriate dilution. Detection of the fusion partner band in the Western blot by serum from the immune rabbit indicates the induction of antibodies specific for fusion partner epitopes.

30 As set forth in Example 10, chimeric particles can be assayed, for example, for induction of cell-mediated immunity specific for fusion peptides, for instance, by injecting chimeric VLPs into mice, and measuring antigen-specific T cell proliferation.

35

As set forth in Example 11, chimeric particles can be assayed, for example, for induction of prophylactic immunity against challenge infection, for instance, by immunizing rabbits with chimeric CRPV VLPs and nonchimeric CRPV VLPs (control), and subsequently challenging with infectious CRPV, to demonstrate that L2 fusions do not abrogate prophylactic immunity; or by immunizing experimental animals with chimeric VLPs containing an STD agent, subsequently challenging with the STD agent, and measuring increased survival against lethal challenge or decreased infection following sub-lethal challenge.

As set forth in Example 12, chimeric particles can be assayed, for example, for induction of therapeutic immunity against pre-existing papillomas, for instance, by immunizing rabbits (that have pre-existing papillomas) using chimeric CRPV VLPs and nonchimeric CRPV VLPs (control), and measuring regression of the pre-existing papillomas.

As set forth in Example 13, chimeric particles can be assayed, for example, for immunotherapy and immunoprevention of tumors.

Particular aspects of the invention may be more readily understood by reference to the following examples, which are intended to exemplify the invention, without limiting its scope to the particular exemplified embodiments.

EXAMPLE 1

Generation of chimeric recombinant baculoviruses

Three chimeras were generated: HPV16L2-HPV16E7, BPVL2-HPV16E7, and BPVL2-HPV16E7 (aa 1-30). HPV16L2-HPV16E7 contained the full-length HPV16E7 fused to the C-terminus (aa 473) of HPV16L2. BPVL2-HPV16E7 contained the full-length HPV16E7 fused to the C-terminus of BPVL2 (aa 469). BPVL2-HPV16E7 (aa 1-30) contained the first 30 amino acids of HPV16E7 fused to the middle of BPVL2 between amino acids 274 and 275.

L2-E7 chimeric genes were generated via recombinant PCR techniques (Higuchi, R. (1990) in PCR Protocols: A Guide to Methods and Applications, ed. Innis, MA, Gelfand, DH, Sninsky, JJ, and White, TJ. (Academic, New York), pp. 177-180). For the chimeras containing full-length HPV16E7, L2 was amplified with a 5' oligo containing a restriction enzyme site, and a 3' oligo which was complementary to the E7 5' oligo. In an independent reaction, E7 was amplified with a 5' oligo complementary to the L2 3' oligo and a 3' oligo containing a restriction enzyme site. The L2 and E7 genes were then fused in a second primer extension reaction using only the outside (L2 5', and E7 3') oligos. For the chimera containing only amino acids 1-30 of HPV16E7, the "internal" primers encoded the first 30 aa of HPV16E7.

The fused genes were then cloned into the baculovirus double expression vector pSynwtVI-, (which already contained L1 cloned under the polyhedrin promoter (Kirnbauer et al., Virology 67:6929 (1993)) immediately downstream of the pSyn promoter.

The BPVL2-HPV16E7 chimeras were cloned as 5' BglII to 3' BglII fragments. The HPV16L2-HPV16E7 chimera was cloned as a 5' SstII to 3' SstII fragment. The primers used for BPVL2-HPV16E7 (full-length) were: BPVL2 sense,

5'GCGGTAGATCTACCTATAAATATGAGTGCACGAAAAAGAGTAAACGT3'(SEQ ID NO:1),

and antisense,

5'GCAATGTAGGTGTATCTCCATGCATGGCATGTTTCCGTTTTTTCGTTTCTCAACAAGGAGGG3'(SEQ ID NO:2),

HPV16E7, sense,

5'CCCTCCTTGTGAGGAAACGAAAAAACGGAAACATGCCATGCATGGAGATACACCTACATTGC3'(SEQ ID NO:3),

and antisense,

5'CCGCTAGATCTGGTACCTGCAGGATCAGCCATGG3'(SEQ ID NO:4).

5 The primers used for BPVL2-HPV16E7 (aa 1-30) were as follows: BPVL2, sense: same as above.

Internal primer, antisense,

5'CAGTTGTCTCTGGTTGCAAATCTAACATATATTCATGCAATGTAGGTGTATCTCCATGCATGGATGAAAACACTTCAGG
ATCTTCCGTGGGC3'(SEQ ID NO:5).

Internal primer, sense,

10 5'GCATGAATATATGTTAGATTGCAACCAGAGACAACCTGATCTCTACTGTTATGAGCAATTAAATGACCAAACATTTCG
AAACCCACTGTATGAAGCAGAACC3'(SEQ ID NO:6).

BPVL2, antisense,

5'CCGCTAGATCTAGGGAGATACAGCTTCTGGCCTTGTGCCACAACGC3'(SEQ ID NO:7).

The primers used for HPV16L2-HPV16E7 chimeras were as follows: HPV16L2, sense,

15 5'GCGGTCCGCGGAATATGCGACACAAACGTTCTGCAAAACGCACAAAACGT3'(SEQ ID NO:8),

and antisense,

5'ATCTCCATGCATGGCAGCCAAAGAGAC3'(SEQ ID NO:9),

HPV16E7, sense,

5'GTCTCTTTGGCTGCCATGCATGGAGAT3'(SEQ ID NO:10),

20 and antisense,

5'GCTCCGCGGGGTACCTGCAGGATCAGCC3'(SEQ ID NO:11).

EXAMPLE 2

Selection of recombinant baculoviruses

25 CsCl-purified recombinant plasmid was cotransfected with baculovirus DNA (Baculo-Gold; PharMingen, San Diego, CA) into Sf9 insect cells (ATCC; CRL 1711) by using Lipofectin (GIBCO/BRL, Gaithersburg, MD) (Hartig, P.C. Biotechniques 11:310 (1991)). The recombinant baculoviruses were plaque-purified as described (Summers, M.D. & Smith, G.E. (1987) A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Texas Agricultural Experiment Station Bulletin (Tex. Agric. Exp. Stn., College Station, TX), Vol 1555).

EXAMPLE 3

Purification of chimeric particles

30 Sf9 insect cells were infected at a multiplicity of infection of 10 with recombinant baculoviruses. After 72 h, the harvested cells were sonicated in phosphate-buffered saline (PBS) for 60 sec. After low-speed clarification, the lysates were subjected to centrifugation at 110,000 x g for 2.5 h through a 40% (wt/vol) sucrose/PBS cushion (SW-28 rotor). The resuspended pellets were centrifuged to equilibrium at 141,000 x g for 20 h at room temperature
35 in a 10-40% (wt/wt) CsCl/PBS gradient. The visible band was harvested, centrifuged to equilibrium again by using the identical conditions, dialyzed extensively against PBS, and stored at 4°C.

- 13 -

EXAMPLE 4**Co-sedimentation of L1/L2-E7 complexes**

5

To determine whether the L2-E7 fusion protein was incorporated into particles, analytical gradient centrifugation was performed as previously described for determining the co-assembly of L1 and L2 (Kirnbauer et al., Virol. 67:6929 (1993)).

10 Briefly, a 12 to 45% sucrose step gradient was allowed to linearize overnight at 4°C, dialyzed samples were layered on top, and the gradient was centrifuged at 41,000 rpm (288,000 x g) for 2 h in a SW-41 rotor. Fractions were harvested. The fractions were analyzed by Coomassie stained SDS-PAGE (L1) or Western blotting with an anti-BPVL2 polyclonal Ab, or anti-HPV16E7 polyclonal Ab, and ¹²⁵I-labeled anti-rabbit IgG.

15 Association of the chimeric L2-E7 with the virus-like particles was established by co-sedimentation.

EXAMPLE 5**Co-immunoprecipitation of L1/L2-E7 complexes**

20 To obtain evidence that the L2-E7 fusion proteins formed stable complexes with L1, co-immunoprecipitation experiments were performed.

Briefly, BPVL1/L2-HPV16E7 (full-length) and BPVL1/L2-HPV16E7 (aa 1-30) VLP preparations were immunoprecipitated in PBS, 1% Triton® X-100 with anti-L1 Mab 5B6 (Roden et al., J. Virol., [supra] (November, 1994)), an anti-L1 polyclonal Ab, 25 pre-immune serum or an irrelevant Ab (anti-E1A Mab) and protein A-Sepharose and subjected to SDS-PAGE. Proteins were immunoblotted and probed with anti-BPVL2, or anti-HPV16E7 sera, or an anti-HPV16E7 Mab (Triton Diagnostics, Alameda, CA).

Association of the chimeric L2-E7 with the virus-like particles was established by co-immunoprecipitation.

- 13a -

EXAMPLE 6

5

Production of antisera

Rabbits were immunized by subcutaneous injection of 330 μ l of CsCl gradient-purified particles in PBS at a concentration of 1 mg/ml. Rabbits were boosted with the same amount of particles two weeks and four weeks after the primary injection.

10

EXAMPLE 7**BPV1 neutralization assay**

A focus-forming assay was performed as described (Kirnbauer et al., Proc. Natl. Acad. Sci. USA 89:12180 (1992)).

Briefly, serial dilutions of rabbit sera obtained 3 weeks after the second booster injection were incubated with \approx 500 focus-forming units of BPV1 virus for 30 min, the virus was adsorbed to C127 cells for 1 h, and the cells were cultured for 3 weeks (Dvoretzky et al., Virology 103:369 (1980)). The foci were stained with 0.5% methylene blue/0.25% carbol fuchsin in methanol.

Equivalent neutralizing titers of 30,000 were obtained for the BPVL1/L2 VLPs and BPVL1/L2-HPV16E7 (full-length) chimeric VLPs.

EXAMPLE 8

Electron microscopy

Transmission electron microscopy was performed as described (Kirnbauer et al., Proc. Natl. Acad. Sci. USA 89:12180 (1992))

5 Briefly, purified particles were adsorbed to carbon-coated grids, stained with 1% uranyl acetate, and examined with a Philips electron microscope model EM 400T at x36,000 magnification.

Particles containing the L2-E7 fusion protein were found to be indistinguishable from L1 or L1/L2 VLPs in terms of morphology and efficiency of incorporation.

EXAMPLE 9

Induction of antibodies

10 Rabbits inoculated with BPVL1/L2-HPV16E7 papillomavirus-like particles produced antisera that recognized E7.

To analyze the sera for E7 specificity, 2.5 μ g of HPV16E7 and 2.5 μ g of BSA (control) were subjected to SDS-PAGE and then Western blotted. Immune and preimmune (control) sera were applied at a dilution of 1:10.

15 Serum from the immune rabbit specifically detected the HPV16E7 protein band in the Western blot, indicating the induction of antibodies specific for E7 epitopes.

Example 10

Induction of cell-mediated immunity

20 Chimeric particles are assayed for induction of cell-mediated immunity specific for E7 peptides, for example, by injecting chimeric VLPs into mice, and measuring antigen-specific T cell proliferation.

Example 11

Prophylactic immunity

25 Chimeric particles are assayed for induction of prophylactic immunity against challenge infection, for example, by immunizing rabbits with E7 chimeric CRPV VLPs and nonchimeric CRPV VLPs (control), and subsequently challenging with infectious CRPV, to demonstrate that L2-E7 fusions do not abrogate prophylactic immunity; or by immunizing experimental animals with chimeric VLPs containing an STD agent, subsequently challenging with the STD agent, and measuring increased survival against lethal challenge or decreased infection following sub-lethal challenge.

Example 12

Therapeutic immunity

30 Chimeric particles are assayed for induction of therapeutic immunity against pre-existing papillomas, for example, by immunizing rabbits (that have pre-existing papillomas) using E7 chimeric CRPV VLPs and nonchimeric CRPV VLPs (control), and measuring regression of the pre-existing papillomas.

Example 13

Immunotherapy and immunoprevention of tumors

35 Chimeric particles are assayed for the ability to prevent tumor development or to treat existing tumors, for example, in experimental animals such as mice using tumor cells expressing, e.g., E7. Animals are immunized with,

- 15 -

e.g., L2-E7 chimeric VLPs and tested for growth of inoculated tumorigenic cells that express, *e.g.*, E7. This approach has been shown to work for animals immunized with
5 noncomplexed E7 and a co-stimulatory protein (Chen et al., Proc. Natl. Acad. Sci. USA 88:110 (1991)).

EXAMPLE 14

Targeting the surface of chimeric VLPs

10

It may be desirable for certain molecules, such as co-stimulators or ligands that bind to cell-surfaces, to be present on the surface of chimeric VLPs. Roden et al., J. Virol., [supra] (November, 1994) determined that regions of L2 are located on the surface of VLPs. A peptide composed of BPV L2 amino acids 44-173, when inoculated into rabbits,
15 induced neutralizing antibodies that were active at a 1:1000 serum dilution. These results indicate that regions of L2 specify neutralizing epitopes, and that these regions are therefore accessible to the surface of native virions. Consequently, fusing a non-L2 polypeptide to one of these regions would likely result in the targeting of this polypeptide to the surface of a chimeric VLP.

20

While particular embodiments of the invention have been described in detail, it will be apparent to those skilled in the art that these embodiments are exemplary rather than limiting, and the true scope of the invention is that defined within the appended claims.

2201601

WO 96/11274

PCT/US95/12914

-16-

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANT: The United States of America

(ii) TITLE OF THE INVENTION: CHIMERIC PAPILLOMA VIRUS-LIKE PARTICLES

(iii) NUMBER OF SEQUENCES: 11

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: Knobbe, Martens, Olson and Bear
- (B) STREET: 620 Newport Center Drive 16th Floor
- (C) CITY: Newport Beach
- (D) STATE: CA
- (E) COUNTRY: USA
- (F) ZIP: 92660

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Diskette
- (B) COMPUTER: IBM Compatible
- (C) OPERATING SYSTEM: DOS
- (D) SOFTWARE: FastSEQ Version 1.5

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Ways Vensko, Nancy
- (B) REGISTRATION NUMBER: 36,298
- (C) REFERENCE/DOCKET NUMBER: NIH032.001CP2

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 619-235-8550
- (B) TELEFAX: 619-235-0176
- (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

.....

GCGGTAGATC TACCTATAAA TATGAGTGCA CGAAAAAGAG TAAAACGT

48

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCAATGTAGG TGTATCTCCA TGCATGGCAT GTTCCGTTT TTTTCGTTTC CTCAACAAGG 60
AGGG 64

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCCTCCTTGT TGAGGAAACG AAAAAAACGG AAACATGCCA TGCATGGAGA TACACCTACA 60
TTGC 64

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCGCTAGATC TGGTACCTGC AGGATCAGCC ATGG

34

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 92 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

2201601

WO 96/11274

PCT/US95/12914

-18-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CAGTTGTCTC TGGTTGCAAA TCTAACATAT ATTCATGCAA TGTAGGTGTA TCTCCATGCA 60
TGGATGAAAA CACTTCAGGA TCTTCCGTGG GC 92

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 102 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCAATGAATAT ATGTTAGATT TGCAACCAGA GACAACTGAT CTCTACTGTT ATGAGCAATT 60
AAATGACCAA ACATTGCAA ACCCACTGTA TGAAGCAGAA CC 102

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCGCTAGATC TAGGGAGATA CAGCTTCTGG CCTTGTTGCC ACAACGC 47

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGGTCCGCG GAATATGCGA CACAAACGTT CTGCAAAACG CACAAAACGT

50

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATCTCCATGC ATGGCAGCCA AAGAGAC

27

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GTCTCTTTGG CTGCCATGCA TGGAGAT

27

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCTCCGCGGG GTACCTGCAG GATCAGCC

28



CA 01201601 1999-04-30

20

WHAT IS CLAIMED IS:

- 5 1. A papillomavirus-like particle having conformational epitopes comprising a papillomavirus L1 fusion product.
2. The papillomavirus-like particle of claim 1, wherein
said papillomavirus L1 fusion product is a human
10 papillomavirus L1 fusion product or a bovine papillomavirus L1 fusion product.
3. The papillomavirus-like particle of claim 2, wherein
said human papillomavirus L1 fusion product is a HPV16 L1
15 fusion product and said bovine papillomavirus L1 fusion product is a BPV1 L1 fusion product.
4. The papillomavirus-like particle of claim 1, wherein
said papillomavirus L1 fusion product comprises a fusion
20 partner that is a peptide or a full-length protein, or comprises fusion partners that include peptides or full-length proteins or combination thereof linked in tandem.
- 25 5. The papillomavirus-like particle of claim 4, wherein said fusion partner is a papillomavirus E6 or E7 product.
6. The papillomavirus-like particle of claim 1, wherein
said papillomavirus L1 fusion product is fused at its
30 N-terminus or C-terminus to a fusion partner.
7. The papillomavirus-like particle of claim 1, wherein
said papillomavirus L1 fusion product has a fusion partner
inserted between L1 amino acids.
- 35 8. The papillomavirus-like particle of claim 1, further comprising a papillomavirus L2 product.





CA 01101601 1999-11-16

21

9. The papillomavirus-like particle of claim 8, wherein said papillomavirus L2 product is a HPV16 L2 product or a BPV1 L2 product.
- 5
10. A composition comprising the papillomavirus-like particle of any one of claims 1-9, and saline.
11. A papillomavirus-like particle characterized as having conformational epitopes,
- 10 comprising a papillomavirus L1 product and a papillomavirus L2 fusion product.
12. The papillomavirus-like particle of claim 11, wherein said papillomavirus L2 fusion product is characterized as being a human papillomavirus L2 fusion product or bovine papillomavirus L2 fusion product.
- 15
13. The papillomavirus-like particle of claim 12, wherein said human papillomavirus L2 fusion product is characterized as being HPV16 L2 fusion product and said bovine papillomavirus L2 fusion product is characterized as being a BPV1 L2 fusion product.
- 20
14. The papillomavirus-like particle of claim 11, wherein said papillomavirus L2 fusion product comprises a fusion partner characterized as being a peptide or a full-length protein, or comprises fusion partners that include peptides or full-length proteins or combinations thereof linked in tandem.
- 25
15. The papillomavirus-like particle of claim 14, wherein said fusion partner is a papillomavirus E6 or E7 product.
16. The papillomavirus-like particle of claim 11, wherein said papillomavirus L2 fusion product is characterized as being fused at its N-terminus or C-terminus to a fusion partner.

30





CA 02101601 1999-11-16

22

17. The papillomavirus-like particle of claim 11, wherein said papillomavirus L2 fusion product is characterized as having a fusion partner inserted between L2 amino acids.

5

18. The papillomavirus-like particle of claim 11, wherein said papillomavirus L1 product is characterized as being a human papillomavirus L1 product or bovine papillomavirus L1 product.

10 19. The papillomavirus-like particle of claim 18, wherein said human papillomavirus L1 product is characterized as being a HPV16 L1 product and said bovine papillomavirus L1 product is characterized as being a BPV1 L1 product.

15 20. The papillomavirus-like particle of claim 11, wherein the particle consists of L1 product and L2 fusion product, and is selected from the group consisting of: (1) HPV16L1 and HPV16L2-HPV16E7 (full-length), wherein HPV16E7 is fused to the C-terminus of HPV16L2, (2) BPV1L1 and BPV1L2-HPV16E7 (full-length), wherein HPV16E7 is fused to the C-terminus of BPV1L2, and (3) BPV1L1 and BPV1L2-HPV16E7 (amino acids 1-30), wherein amino acids 1-30 of HPV16E7 are fused to BPV1L2 between amino acids 274 and 275.

20

21. A method of purification of the papillomavirus-like particle of claim 11, comprising the step of exposing the papillomavirus-like particle of claim 11, to an affinity chromatography column, comprising antibodies that bind to a fusion partner of the papillomavirus L2 fusion product of the papillomavirus-like particle of claim 11, resulting in
25 the purification of said particle.

22. A method of purification of a fusion partner of the papillomavirus L2 fusion product of the papillomavirus-like particle of claim 11, comprising the step of isolating the papillomavirus-like particle of claim 11 resulting in the purification of said fusion partner.

30





CA 02201601 1999-11-16

23

23. Use of the papillomavirus-like particle of any one of claims 11-20 for delivery into a cell of a fusion partner of the papillomavirus L2 fusion product.

5

24. An immunogenic composition comprising the papillomavirus-like particle of any one of claims 11-20, and saline.

25. The use of a papillomavirus-like particle of claim 11 to stimulate an immune response in a host.

10

26. A synthetic DNA molecule characterized as encoding a papillomavirus L1 product and papillomavirus L2 fusion product, wherein said molecule is capable of directing expression in a transformed cell of papillomavirus-like particles characterized as having conformational epitopes, wherein said particles comprise said papillomavirus L1 product and papillomavirus L2 fusion product.

15

27. The DNA molecule of claim 26, wherein the DNA molecule is capable of directing expression in a transformed insect cell and the DNA further comprises an insect cell vector, or wherein the DNA molecule is capable of directing expression in a transformed mammalian cell and the DNA further comprises a mammalian cell vector, or wherein the DNA molecule is capable of directing expression in a transformed yeast cell and the DNA further comprises a yeast cell vector.

20

28. A host cell transformed with one or more DNA molecules which together encode a papillomavirus L1 product and a papillomavirus L2 fusion product, wherein said molecule(s) direct(s) expression of papillomavirus-like particles characterized as having conformational epitopes, wherein

25

30





CA 01201601 1999-04-30

24

said particles comprise said papillomavirus L1 product and papillomavirus L2 fusion product.

5

29. The host cell of claim 28, wherein the papillomavirus L1 product and the papillomavirus L2 fusion product are encoded on the same DNA molecule.

10

30. The host cell of claim 28, wherein the papillomavirus L1 product and the papillomavirus L2 fusion product are encoded on different DNA molecules.

15

31. A method for producing a papillomavirus-like particle, characterized as having conformational epitopes, comprising a papillomavirus L1 product and a papillomavirus L2 fusion product, which method comprises providing conditions for the host cell of claim 28 to express said DNA, thereby producing said particle.

20

32. The method of claim 31 further comprising recovering said particle from the host cell.

25

33. A composition comprising a first synthetic DNA molecule characterized as encoding a papillomavirus L1 product and a second synthetic DNA molecule characterized as encoding a papillomavirus L2 fusion product, wherein said molecules are capable of directing expression in a transformed cell of papillomavirus-like particles characterized as having conformational epitopes, wherein said particles comprise said papillomavirus L1 product and papillomavirus L2 fusion product.

30

34. The composition of claim 33, wherein the DNA molecules are capable of directing expression in a transformed insect cell and the DNA of each said molecule further comprises an insect cell vector, or wherein the DNA molecules are capable of directing expression in a transformed mammalian cell and the DNA of each said molecule further comprises a

35





CA 01101601 1999-04-30

25

- 5 mammalian cell vector, or wherein the DNA molecules are capable of directing expression in a transformed yeast cell and the DNA of each said molecule further comprises a yeast cell vector.

